

# Microscopics of meson degrees of freedom in nucleons and mesons in nuclei - what can be seen in the process of quasielastic knockout of mesons by high-energy electrons

V.G. Neudatchin, I.T. Obukhovskiy, L.L. Sviridova, N.P.Yudin\*

D.V. Skobeltsyn Institute of Nuclear Physics, Moscow State University 119899 Moscow, Russia

## Abstract

Developed earlier concept of quasielastic knock out of pions from nucleons by high- energy electrons is propounded as a tool for checking microscopical model (  $^3P_0$  - fluctuation ) for decay of  $N$  to different channels  $\pi + B$  and Preparata model of nucleus structure.

## 1 Introduction

1. The quasielastic knock-out (q.e.k.) of protons out of nuclei by high energy electrons was used as mighty tool for investigation of nucleus structure [1]. Then the concept of q.e.k. was extended [2] to the knock-out of electrons out of atoms and molecules and by this method very important results for the structure of these microobjects were obtained. For recent time in the literature, the notion of quasielastic knock out is often discussed in hadron physics [3]. In general, this term means that in a process almost whole four momentum of a probe is transferred to only one degree of freedom - for instance to nucleon in the process of photoproduction of pions on nuclei.

We have extended this notion to the quasielastic knock out of mesons from nucleon and nuclei and use this notion to clarify some physical problems. In general, the set of diagrams gives contributions to the process of photo- or electroproduction of pions on nucleons. For example, in Fig. 1 we show several such diagrams (with pseudoscalar interaction). In the case of photoproduction all of these diagrams play very important role. However at sufficiently high square momentum transfer  $Q^2$  ( $Q^2 \geq 1 \text{ (GeV/c)}^2$ ), and total invariant mass of hadrons  $W \geq 2 \text{ GeV/c}$  (in order to exclude the contributions of resonances) the situation becomes very simple - for longitudinal photons the "pion in flight diagram", that is the diagram with pion pole in  $t$ -channel, becomes dominant. This was shown in our previous papers [4]. It means that in the longitudinal electroproduction in the mentioned kinematics the real knock-out of pion from nucleon takes place. Thus from experimental data one can extract the momentum distribution of pion in nucleon and their spectroscopic factors i.e. the number of pions. This notion opens a new possibility for direct investigation of meson structure of nucleon and relevant problems. The similar considerations are also valid for

---

\*e-mail:yudin@helena.sinp.msu.ru




Figure 1: The Born diagrams of photo- and electro-production of pions on nucleon.

electroproduction of pions by transverse virtual photons and we have here the knock-out of vector mesons with simultaneous transformation of them into pion (the phenomenon of "deexcitation" that is similar to the deexcitation of alpha - particles in nuclei). It can also provide a valuable information on the dynamic of vector mesons in nucleon.

2. Now let's make several remarks about terms in which we shall discuss these problems. The cross section of electroproduction has the following form:

$$\frac{d\sigma(\gamma^*p \rightarrow n\pi^+)}{dWdt} \sim \varepsilon\sigma_L + \sigma_T \quad (1)$$

where  $\varepsilon$  is a usual polarization parameter of the virtual photon,  $t$  - momentum square transferred to nucleon,  $\sigma_L$ ,  $\sigma_T$  - longitudinal and transverse cross sections. Analytically the hadron part of pion in flight diagram of Fig.1 is written as follows:

$$\frac{M(p \rightarrow n\pi^+)}{k^2 - m_\pi^2} = \frac{M(p \rightarrow n\pi^+)}{k_0 - e_k} \cdot \frac{1}{k_0 + e_k} = \langle n | a_{\pi^+} | p \rangle \frac{1}{k_0 + e_k} \quad (2)$$

where  $a_{\pi^+}$  is an operator of virtual pion absorption,  $k_0$  - zero component of transferred from nucleon to photon four vector,  $e_k$  - energy of the virtual pion with momentum  $k$  in the rest frame of proton,  $p$  and  $n$  - proton and neutron,  $M(p \rightarrow n\pi^+)$  - invariant amplitude of transformation  $M(p \rightarrow n\pi^+)$ . It is important to point out that this equation takes into account simultaneously the diagrams going "forward and backward" in time, and that we work in the lab system. The quantity  $\langle n | a_{\pi^+} | p \rangle$  defines the momentum distribution of pions in proton and we call it "wave function" and designate by  $\Psi_p^{n\pi}$ . This wave function is normalized by the condition:

$$\int d\tau |\Psi_p^{n\pi}(k)|^2 = S_p^{n\pi} \quad (3)$$

where  $d\tau = d^3k / ((4\pi)^3 E_p E_n)$ ,  $E_{p,n}$  - energies of proton and neutron,  $S_p^{n\pi}$  is a spectroscopic factor of pion in nucleon. It determines the number of pions participating in production of the channel. The sum of the spectroscopic factors over all final channels will give the total number of positive pions in nucleon.

As a result, we get that the longitudinal cross section of pion electroproduction has the following form:

$$\frac{d\sigma_L(\gamma^* p \rightarrow n\pi^+)}{dt} \sim |\Psi_p^{n\pi}(k)|^2 F_{\gamma\pi\pi}^2(Q^2) \quad (4)$$

where  $t = k^2$  and  $F_{\gamma\pi\pi}^2(Q^2)$  - electromagnetic formfactor of pion,  $Q^2$  - "mass" of the virtual photon.

## 2 Check of $^3P_0$ fluctuation model

It is well known, that the hadrons are effective degrees of freedom and the hadron dynamics should be a phenomenological one. The important ingredient of this phenomenology is the coupling constants of meson with baryons. Now when an "attack" on hadrons world is wide underway the number of constant becomes big and it is necessary to search for notions, that can unite the description of hadrons. It is naturally, that the search for unification should be based on quark-gluon degrees of freedom. One way to realize this unification is to introduce the coupling between quarks and mesons. Then the coupling with baryons comes out as a result of some averaging over quark structure of baryons. Partially this was done in the paper [5]. But this approach entails also many parameters. It is proved that an old model of  $^3P_0$  fluctuation has definite advantages. On diagram level this model is presented in Fig.2.




Figure 2: The diagram of vacuum production of the flavour SU3-invariant quark-antiquark pair.

The advantage of this approach based on the fact that the five quarks in the final system can be transformed into hadron by many different ways. This permits one on the same footing to consider and therefore to relate the whole set of meson-baryons coupling constants, for instance pion-baryon, kaon-baryon, vector meson-baryon and so on. The formal aspects of this rearrangement were considered in our papers [6, 7] and will not be discussed here. Calculated momentum distributions and formfactors of pions and kaons in nucleon are shown in Fig.3,4.

One can see that discussed momentum distributions are very interesting and they demand further investigation.

In conclusion we would like to remark that  $^3P_0$ -fluctuation is probably closely related with scalar quark condensate. This is one more reason to investigate this model.




Figure 3: The momentum distributions of pions  $|\Psi_p^{B\pi}|^2$  in different channels ( $\pi B = \pi N, \pi \Delta$ ).




Figure 4: The momentum distributions of pions  $|\Psi_p^{B\pi}|^2$  in different channels ( $B = N^*, N^{**}$ ).  $N^* = N(1535)$ ,  $N^{**} = N(1440)$ .

### 3 Mesons in nuclei

Over past years, the problem of pions in nuclei was discussed in literature [8] many times. For instance, a very well known paper has a title "Where is the pions in nuclei" [9]. We can not dwell on the details of this very interesting questions and confine ourselves with the statement that now the problem is partially clear - in any case, the pions in nuclei predicted by RPA approach were detected and sharp contradiction between prediction of the theory and (p,n) experiment, that was a base for the title of paper [9], was probably removed. In a sense, one really sees the pions in nuclei injected there by RPA correlations [10]. New nontrivial aspects of the nuclear pion problem were considered in the paper [11].

Nevertheless the problem of mesons in nuclei is far from its full solution and probably that we will meet here the unexpected and nontrivial phenomena.

The first modern nontrivial notion of meson's injection into nuclei was propounded by Migdal [12]. Now it seems clear that his notion of pion condensate in nuclei is not realized. But there exists another concept of pion condensate propounded by Preparata [13]. Preparata's idea is closely related to the effect of Dicke [14] that is known in atomic physics for several tens of years.

The point is: let's suppose that one has a system of  $N$  excited atoms. Then as long as radioactive transitions take place the probability of transitions becomes bigger and bigger and in the "middle" - i.e. when about one half of excited atoms is left - the probability becomes as much as  $N^2/4$ . At large  $N$  the increase of probability becomes fantastic! Preparata has assumed that in nuclei something similar can take place if one takes into account  $\Delta$ -resonance. In this case, one has  $\Delta \rightarrow$  excited atoms,  $N \rightarrow$  atom in the ground state, pion  $\rightarrow$  photon. According to an estimate by Preparata if one takes into account the existence of such transitions the increasing of self-energy of whole nucleus will result in gain of energy that amounts 60 MeV per one nucleon! Thus, a heavy nucleus can consist of three coherently vibrating subsystems  $\Delta$ -nucleons,  $\Delta$ -resonances and pions. With the usual way of investigation it is not simple task to observe such a substructures. We suggest

to use for this purpose the quasielastic knock out of pions. In Fig.5,




Figure 5: Momentum distributions of pions in nuclei: solid line - the MD  $|\Psi_p^{n\pi}(k)|^2$ ,  $(\text{GeV}/c)^{-3}$  versus  $|k|$ ,  $\text{GeV}/c$  in Preparata model; dashed line - the MD in the model of independent nucleons.

it is shown the momentum distribution of pions that are injected into nuclei by Preparata mechanism in comparison with trivial injection by independent nucleons. We see that the momentum distributions of pions, injected by usual mechanisms (independent nucleons) and Preparata mechanism are drastically different and measurement of pion momentum distribution helps to prove whether there exists a collective strengthening of pions in nuclei.

## References

- [1] D.G. Ireland and G.Van der Steenhoven Phys.Rev. C49,2182(1994); D.Debruyne, J.Ryckebush et al Phys.Rev.C62, 024611 (2000).
- [2] V.G.Neudatchin, Yu.V. Popov and Yu.F. Smirnov Usp.Fiz.Nauk 169,1111(1999); E.Weigold and McCarthy Electron Momentum Spectroscopy, Plenum, New York,1999.
- [3] X.Li and L.E.Wright, C.Bennhold Phys.Rev.C48,816 (1993).
- [4] V.G.Neudatchin, N.P.Yudin and L.L.Sviridova Yad.Fiz. 64,1680 (2001)
- [5] G.Brown and D.Riska Nucl.Phys.A679 (2001),577
- [6] I.T.Obukhovsky, V.G.Neudatchin, L.L. Sviridova, N.P. Yudin Yad.Fiz. 66(2003)
- [7] I.T.Obukhovsky, V.G.Neudatchin, L.L.Sviridova Yad.Fiz. to be published
- [8] D.S.Koltun Phys.Rev. C57,1210 (1997)
- [9] G.E.Brown, M,Bubala,Zi.Bang, J.Wambach Nucl.Phys.A593(1995)293.

- [10] H.Sakai Nucl.Phys.A690(2001)66c.
- [11] G.Chanfray,M.Ericson and P.A.M.Guichon Phys.Rev. C63(2001)055202.
- [12] A.B. Migdal Rev.Mod.Phys. 50,107(1978).
- [13] G.Preparata Nuovo Cimento A103(1990)1213.R.Alzetta,G. Liberti and G.Preparata  
Nuovo Cimento A112(1999)1609
- [14] A.V. Andreev,V.I. Emel'yanov and L.L. Il'inskii Coherent Phenomena in Optics  
(Nauka,Moscow,1988).